

This is a repository copy of *Khovanov homotopy types and the Dold-Thom functor*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/90031/>

Version: Submitted Version

---

**Article:**

Everitt, Brent [orcid.org/0000-0002-0395-338X](https://orcid.org/0000-0002-0395-338X), Lipshitz, Robert, Sarkar, Sucharit et al. (1 more author) (2016) Khovanov homotopy types and the Dold-Thom functor. *Homology, Homotopy and Applications*. pp. 177-181. ISSN 1532-0081

<https://doi.org/10.4310/HHA.2016.v18.n2.a9>

---

**Reuse**

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.

# Khovanov homotopy types and the Dold-Thom functor

Brent Everitt, Robert Lipshitz, Sucharit Sarkar and Paul Turner<sup>\*</sup>

**Abstract.** We show that the spectrum constructed by Everitt and Turner as a possible Khovanov homotopy type is a product of Eilenberg-MacLane spaces and is thus determined by Khovanov homology. By using the Dold-Thom functor it can therefore be obtained from the Khovanov homotopy type constructed by Lipshitz and Sarkar.

A *Khovanov homotopy type* is a way of associating a (stable) space to each link  $L$  so that the classical invariants of the space yield the Khovanov homology of  $L$ . There are two recent constructions of Khovanov homotopy types, using different techniques and giving different results [3, 6]. In [3] homotopy limits were employed to build an  $\Omega$ -spectrum  $\mathbf{X}_\bullet L = \{X_k(L)\}$  with the following properties:

- (i). the homotopy type is a link invariant, and
- (ii). the homotopy groups are Khovanov homology:

$$\pi_i(\mathbf{X}_\bullet(L)) = Kh^{-i}(L).$$

The main goal of this note is to prove the following result.

**Theorem 1.** *Each of the spaces  $X_k(L)$  is homotopy equivalent to a product of Eilenberg-MacLane spaces.*

In [6] the programme of Cohen, Jones and Segal [2] was generalized to produce a suspension spectrum  $\mathcal{X}_{Kh}(L)$  with the following properties:

- (i). the homotopy type is a link invariant, and
- (ii). the reduced cohomology is Khovanov homology:

$$\tilde{H}^i(\mathcal{X}_{Kh}(L)) = Kh^i(L).$$

As a corollary we obtain that  $\mathbf{X}_\bullet(L)$  is homotopy equivalent to the infinite symmetric product of  $\mathcal{X}_{Kh}(L)$ .

To prove Theorem 1 we use the explicit model, due to McCord [8], of the Eilenberg-MacLane spaces. Given a monoid  $G$  and a based topological space  $X$ , let  $B(G, X)$  denote the set of maps

---

BRENT EVERITT: Department of Mathematics, University of York, York YO10 5DD, United Kingdom. e-mail: [brent.everitt@york.ac.uk](mailto:brent.everitt@york.ac.uk). ROBERT LIPSHITZ, SUCHARIT SARKAR: Department of Mathematics, Columbia University, New York NY 10027, USA. e-mail: [lipshitz@math.columbia.edu](mailto:lipshitz@math.columbia.edu), e-mail: [sucharit@math.columbia.edu](mailto:sucharit@math.columbia.edu). PAUL TURNER: Section de mathématiques, Université de Genève, 2-4 rue du Lièvre, CH-1211, Geneva and Département de mathématiques, Université de Fribourg, Chemin du musée, CH-1700 Fribourg, Switzerland. e-mail: [pri.maths@gmail.com](mailto:pri.maths@gmail.com).

<sup>\*</sup> RL was supported by an NSF grant number DMS-0905796 and a Sloan Research Fellowship. SS was supported by a Clay Foundation Postdoctoral Fellowship. PT was partially supported by Swiss National Science Foundation grant 200021-131967.

$u: X \rightarrow G$  such that  $u(x) = 0$  for all but finitely many  $x \in X$ . Then  $B(G, X)$  is a monoid, and if  $G$  is a group (the case of interest) then  $B(G, X)$  is a group. Moreover, when  $G$  is an abelian topological group the set  $B(G, X)$  can be topologized in a natural way so that the group operation is continuous. This construction has nice functoriality: letting  $\mathbf{Ab}$ ,  $\mathbf{Top}_*$  and  $\mathbf{AbTop}$  denote respectively the categories of abelian groups, based topological spaces and topological abelian groups, one has the following result.

**Proposition 1.** [8, Proposition 6.7] *McCord's construction is a bifunctor*

$$B(-, -): \mathbf{Ab} \times \mathbf{Top}_* \rightarrow \mathbf{AbTop}.$$

Furthermore, as special case of [8, Theorem 11.4], for an abelian group  $G$  the space  $B(G, S^n)$  is the Eilenberg-MacLane space  $K(G, n)$ . Thus we may take as *the* Eilenberg-MacLane space functor:

$$B(-, S^n): \mathbf{Ab} \rightarrow \mathbf{AbTop}.$$

Conversely, the following is [4, Corollary 4K.7, p. 483] (apparently originally due to Moore; cf. [8, p. 295]):

**Proposition 2.** *A path-connected, commutative topological monoid is a product of Eilenberg-MacLane spaces.*

The spaces  $X_k(L)$  are built as homotopy limits of diagrams of spaces. Recall that given a small category  $\mathbf{C}$  and a (covariant) functor  $D: \mathbf{C} \rightarrow \mathbf{Top}_*$  (a diagram), that  $\text{holim}_{\mathbf{C}} D$  is constructed as follows (see, e.g., [1, Section 11.5] or the concise notes [9, Section 3.7]). Consider the product

$$\prod_{\sigma \in N(\mathbf{C})} \text{Hom}(\Delta^n, D(c_n)) = \prod_{n \geq 0} \prod_{\substack{c_0 \xrightarrow{\alpha_1} \dots \xrightarrow{\alpha_n} c_n \\ \alpha_i \neq \text{Id}}} \text{Hom}(\Delta^n, D(c_n)) \quad (1)$$

where  $N(\mathbf{C})$  is the subset of the nerve of  $\mathbf{C}$  consisting of all sequences of composable morphisms  $\sigma = (c_0 \xrightarrow{\alpha_1} c_1 \xrightarrow{\alpha_2} \dots \xrightarrow{\alpha_n} c_n)$  in which none of the morphisms are identity maps, and  $\text{Hom}$  denotes the space of continuous maps from the standard  $n$ -simplex. The homotopy limit  $\text{holim}_{\mathbf{C}} D$  is the subspace of this product consisting of those tuples  $(f_\sigma)_{\sigma \in N(\mathbf{C})}$  such that the following diagrams commute:

$$\begin{array}{ccc} \Delta^{n-1} & \xrightarrow{f_{d_i \sigma}} & D(c_n) \\ d^i \downarrow & & \downarrow \text{Id} \\ \Delta^n & \xrightarrow{f_\sigma} & D(c_n) \end{array} \quad (2)$$

for each  $0 < i < n$ , and

$$\begin{array}{ccc} \Delta^{n-1} & \xrightarrow{f_{d_0 \sigma}} & D(c_n) \\ d^0 \downarrow & & \downarrow \text{Id} \\ \Delta^n & \xrightarrow{f_\sigma} & D(c_n) \end{array} \quad \text{and} \quad \begin{array}{ccc} \Delta^{n-1} & \xrightarrow{f_{d_n \sigma}} & D(c_{n-1}) \\ d^n \downarrow & & \downarrow D(\alpha_n) \\ \Delta^n & \xrightarrow{f_\sigma} & D(c_n) \end{array} \quad (3)$$

corresponding to the cases  $i = 0$  and  $i = n$ , respectively. Here the map  $d^i$  denotes the  $i^{\text{th}}$  face inclusion,  $d_i \sigma = (c_0 \xrightarrow{\alpha_1} \dots \xrightarrow{\alpha_{i-1}} c_{i-1} \xrightarrow{\alpha_{i+1} \alpha_i} c_{i+1} \dots \xrightarrow{\alpha_n} c_n)$  when  $0 < i < n$ , and  $d_0, d_n$  similarly.

The following is well-known, but for completeness we give its (short) proof.

**Proposition 3.** *Let  $D: \mathcal{C} \rightarrow \text{Top}_*$  be a diagram of topological abelian groups and continuous group homomorphisms. Then the homotopy limit of  $D$  is a topological abelian group.*

*Proof.* Pointwise addition makes the set  $\text{Hom}(\Delta^n, D(c_n))$  into an abelian group, and the product in formula (1) is the product (topological abelian) group. It remains to see that the diagrams (2) and (3) describe a subgroup of this product. Suppose that tuples  $(f_\sigma)$  and  $(g_\sigma)$  make these diagrams commute. Then the first two diagrams automatically commute for the pointwise sum  $(f_\sigma + g_\sigma)$ . The third diagram for the pointwise sum becomes,

$$\begin{array}{ccccccc} \Delta^{n-1} & \longrightarrow & \Delta^{n-1} \times \Delta^{n-1} & \xrightarrow{f_{d_n\sigma} \times g_{d_n\sigma}} & D(c_{n-1}) \times D(c_{n-1}) & \xrightarrow{+} & D(c_{n-1}) \\ d^n \downarrow & & \downarrow d^n \times d^n & & \downarrow D(\alpha_n) \times D(\alpha_n) & & \downarrow D(\alpha_n) \\ \Delta^n & \longrightarrow & \Delta^n \times \Delta^n & \xrightarrow{f_\sigma \times g_\sigma} & D(c_n) \times D(c_n) & \xrightarrow{+} & D(c_n) \end{array}$$

for which the first square obviously commutes, the second commutes since  $f$  and  $g$  are in the prescribed subspace and the third commutes from the fact that  $D(\alpha_n)$  is a group homomorphism. The inverse operation is similarly seen to be closed, hence the subspace defined above is a subgroup.  $\square$

*Proof (Proof of Theorem 1).* Let  $L$  be an oriented link diagram with  $c$  negative crossings. The space  $X_k(L)$  is constructed as follows. Let  $I$  denote the category with objects  $\{0, 1\}$  and a single morphism from 0 to 1, and  $I^n$  the product of  $I$  with itself  $n$  times. Let  $\bar{0}$  be the initial object in  $I^n$ , and let  $\mathbf{P}$  be the result of adjoining one more object to  $I^n$  and a single morphism from the new object to every object except  $\bar{0}$ .

In [3] it is shown that there is a functor  $F: \mathbf{P} \rightarrow \text{Ab}$  such that the  $i^{\text{th}}$  derived functor of the inverse limit,  $\varprojlim_{\mathbf{P}}^i F$ , is isomorphic to the  $i^{\text{th}}$  unreduced Khovanov homology of  $L$ . The space  $X_k(L)$  is constructed by composing this functor with the Eilenberg-MacLane space functor  $K(-, k+c)$  and taking the homotopy limit of the resulting diagram of spaces.

We may now use the explicit model for Eilenberg-MacLane spaces given by McCord. By applying Proposition 1 we define a diagram  $D: \mathbf{P} \rightarrow \text{AbTop}$  as the composition

$$\mathbf{P} \xrightarrow{F} \text{Ab} \xrightarrow{B(-, S^{k+c})} \text{AbTop}.$$

By the homotopy invariance property of the homotopy limit construction we have

$$X_k(L) \simeq \text{holim}_{\mathbf{P}} D.$$

By Proposition 3, the homotopy limit on the right is itself a topological abelian group, and hence, by Proposition 2, a product of Eilenberg-MacLane spaces.  $\square$

**Corollary 1.** *The homotopy type of  $\mathbf{X}_\bullet(L)$  is determined by  $Kh(L)$ .*

The spectrum  $\mathcal{X}_{Kh}(L) = \{\mathcal{X}_{Kh}^{(k)}(L)\}$  constructed in [6] has the additional property that the cellular cochain complex of the space  $\mathcal{X}_{Kh}^{(k)}(L)$  is isomorphic to the Khovanov complex of  $L$  (up to shift). It follows from the description of the Khovanov homology of the mirror image (see [5]) that

$$\tilde{H}_i(\mathcal{X}_{Kh}(L)) = Kh^{-i}(-L)$$

where  $-L$  denotes the mirror of  $L$ . The infinite symmetric product  $\mathrm{Sym}^\infty \mathcal{X}_{Kh}^{(k)}(L)$  is seen from the Dold-Thom theorem to be

$$\mathrm{Sym}^\infty \mathcal{X}_{Kh}^{(k)}(L) = \prod_n K(\tilde{H}_n(\mathcal{X}_{Kh}^{(k)}(L)), n)$$

from which we have the following.

**Corollary 2.** *For large enough  $k$ , the space  $X_k(-L)$  is homotopy equivalent to the infinite symmetric product  $\mathrm{Sym}^\infty \mathcal{X}_{Kh}^{(k)}(L)$ .*

We end by noting that the analogue of Theorem 1 for the spectra  $\mathcal{X}_{Kh}(L)$  is not true. For all alternating knots  $\mathcal{X}_{Kh}(L)$  is a wedge of Moore spaces [6], however there are examples of non-alternating knots for which  $\mathcal{X}_{Kh}(L)$  is not a wedge of Moore spaces (see [7]).

### Acknowledgements

We thank Tyler Lawson for several helpful suggestions, including communicating Proposition 3 to us.

### References

- [1] A. K. Bousfield and D. M. Kan, *Homotopy limits, completions and localizations*, Lecture Notes in Mathematics, Vol. 304, Springer-Verlag, Berlin, 1972. MR0365573 (51 #1825)
- [2] R. L. Cohen, J. D. S. Jones, and G. B. Segal, *Floer's infinite-dimensional Morse theory and homotopy theory*, The Floer memorial volume, 1995, pp. 297–325. MR1362832 (96i:55012)
- [3] Brent Everitt and Paul Turner, *The homotopy theory of Khovanov homology*. arXiv:1112.3460.
- [4] Allen Hatcher, *Algebraic topology*, Cambridge University Press, Cambridge, 2002. MR1867354 (2002k:55001)
- [5] Mikhail Khovanov, *A categorification of the Jones polynomial*, Duke Math. J. **101** (2000), no. 3, 359–426. MR1740682 (2002j:57025)
- [6] Robert Lipshitz and Sucharit Sarkar, *A Khovanov homotopy type or two*. arXiv:1112.3932.
- [7] ———, *Some Steenrod squares on Khovanov homology*. In preparation.
- [8] M. C. McCord, *Classifying spaces and infinite symmetric products*, Trans. Amer. Math. Soc. **146** (1969), 273–298. MR0251719 (40 #4946)
- [9] Rubén Sánchez-García, *Homotopy limits and colimits*. [www.math.uni-duesseldorf.de/~sanchez/](http://www.math.uni-duesseldorf.de/~sanchez/).